



- (72) LUBITZ, Werner, AT
(72) JECHLINGER, Wolfgang, AT
(72) SZOSTAK, Michael, AT
(72) WITTE, Angela, AT
(71) EVAX TECHNOLOGIES GMBH VACCINE DEVELOPMENT, DE
(51) Int.Cl.⁶ C12N 15/73, C12Q 1/68, C12N 1/21, C12N 15/10, A61K 39/02
(30) 1996/08/21 (196 33 698.8) DE
(54) **NOUVEAUX SYSTEMES DE REGULATION DE L'EXPRESSION
GENETIQUE**
(54) **NEW SYSTEMS FOR REGULATING GENETIC EXPRESSION**

(57) La présente invention concerne un procédé pour sélectionner de nouvelles séquences d'opérateurs P_R ou P_L constituées de phages lambdoïdes qui présentent une thermostabilité différente de celle des séquences de type sauvage, pour la fixation d'un répresseur. L'invention concerne en outre de nouvelles séquences d'opérateurs P_R ou P_L mutées, ainsi que leur utilisation pour l'expression génétique thermorégulée et pour la préparation de vaccins améliorés.

(57) The present invention concerns a process for selecting new P_R - or P_L -operator sequences of lambdoid phages which, compared to wild-type sequences, have a different thermostability for the binding of a repressor. In addition, the invention discloses new mutated P_R - or P_L -operator sequences and their use for temperature-regulated expression of genes and for producing improved vaccines.



- 1 -

New systems for the regulation of gene expression**Description**

The present invention concerns a method for selecting new P_R or P_L operator sequences from lambdoid phages which have a different thermostability compared to the wild-type sequence with regard to binding a repressor. In addition new mutated P_R or P_L operator sequences and their application for the temperature-regulated expression of genes and for the production of improved vaccines are disclosed.

The initiation of transcription of the O_R - O_L region of the bacteriophage lambda and other lambdoid phages is negatively and positively regulated by a repressor which is the product of the cI gene (see review article Ptashne et al., Cell 19 (1980), 1-11). In the O_R region three operator sequences (O_{R1} , O_{R2} and O_{R3}) overlap the promoters P_R and P_{RM} which are orientated in different directions. P_R controls the transcription of genes which are responsible for the lytic multiplication cycle of the phage whereas P_{RM} is the promoter for the lambda cI gene which is responsible for maintaining the lysogenic state. The lambda repressor cI binds co-operatively to the operator sequences O_{R1} and O_{R2} with the result that P_R is repressed and P_{RM} is activated.

In addition the bacteriophage lambda also contains a further operator region O_L which also contains three operator sequences (O_{L1} , O_{L2} and O_{L3}). The expression of the lambda N gene can be repressed by the P_L promoter by binding of the cI repressor to this O_L operator region.

- 2 -

Promoters of the bacteriophage lambda in particular the P_L and the P_R promoter have been used for a long time in recombinant DNA technology for heterologous temperature-regulated gene expression in *E. coli* (cf. Hedgpeth et al., *Molec. Gen. Genet.* 183 (1978), 197-203 and Bernard et al., *Gene* 5 (1979), 59-76; Buell et al., *Nucleic Acids Res.* 13 (1985), 1923 and Shatzman and Rosenberg, *Methods Enzymol.* 152 (1987), 661). A temperature-sensitive lambda repressor $cI857$ is used in these expression systems which represses the P_L and P_R transcription at low temperatures up to 30°C but allows a gene expression at higher temperatures.

An advantage of this lambda expression system is that the gene expression can be induced in a simple manner by increasing the temperature and no addition of chemical inducers is necessary for this. However, a serious disadvantage is that the repression of gene expression only occurs up to relatively low temperatures of not more than 30°C, which is a temperature at which only a slow bacterial growth occurs. Hence the object of the invention was to provide an improved system for lambda P_L or P_R gene expression which enables a repression at variable higher temperatures.

This object is achieved by providing mutated P_R or P_L operator sequences from lambdoid phages which, compared to the wild-type operator sequence, have a different and in particular higher thermostability with regard to the binding of a temperature-sensitive repressor. The finding that lambda expression systems with an improved thermostability can be produced at all is extremely surprising since, apart from the temperature-sensitive lambda $cI857$ mutant, no other temperature-sensitive cI mutants are known but only those mutations in the cI

- 3 -

repressor are known which make the molecule more resistant to thermal inactivation (Hecht et al., Proteins 1 (1986), 43-46 and Das and Mandal, Mol.Gen.Genet. 204 (1986), 540-542). It was even more surprising that mutations which lead to an improved thermostability are located in the operator DNA sequence and not in the DNA sequence coding for the repressor molecule. Thus for example a mutation of the lambda O_R2 operator sequence is known from the literature which leads to a complete loss of repressor binding (Hawley et al., J.Biol.Chem. 260 (1985), 8618-8626).

A method is provided for identifying suitable mutants which enables the selection of mutated O_R or O_L operator DNA sequences from lambdoid phages which have a different thermostability compared to the wild-type sequence with regard to binding a repressor in which the method is characterized in that (a) a DNA cassette is prepared which contains a selection gene under the operative control of an expression control sequence comprising at least one O_R or O_L operator sequence from a lambdoid phage and a promoter, (b) the operator DNA sequence is subjected to a mutagenesis and (c) the mutated operator DNA sequences are analysed.

The lambdoid phages are preferably selected from the group comprising the phage lambda, phage 21, phage 22, phage 82, phage 424, phage 434, phage D326, phage DLP12, phage gamma, phage HK022, phage P4, phage Phi80, phage Phi81, coliphage 186 and recombinant variants thereof. The said phages are very similar with regard to the mechanism of repression of gene expression by means of a cI repressor (Johnson et al., Nature 294 (1982), 217-223). Recombinant variants of the said phages e.g. lambda imm434 can be obtained by substitution of

- 4 -

individual genome fragments within the said phages (cf. for this Hendricks et al., Lambda 2 (1983), R.W. Hendricks, J.W. Roberts, F.W. Stahl and R.A. Weissberg (publisher), Cold Spring Harbor Laboratory Press, New York). The phage lambda or a recombinant variant thereof is preferably used as the lambdoid phage e.g. lambda imm434. An operator DNA sequence from the operator regions O_R (SEQ ID NO.1) or /and O_L (SEQ ID NO.3) of the phage lambda and in particular one of the operator sequences O_{R1} , O_{R2} and O_{R3} or O_{L1} , O_{L2} and O_{L3} contained therein is particularly preferably used for the mutagenesis. The operator sequence O_{R2} is most preferred.

The selection gene for the DNA cassette which is brought under the operative control of the expression control sequence containing the mutated operator sequence, preferably a lambda operator/promoter region, is preferably a suicide gene which when expressed leads to the death of the bacterial cell and thus serves as a selection marker for identifying suitable mutants. The suicide gene should be so strongly repressed at a temperature at which the lambda repressor binds to the mutated operator sequence that a bacterial cell containing the DNA cassette can grow. When the maximum temperature at which the repressor can still bind to the operator is exceeded, the suicide gene is expressed and the bacterial cell is destroyed. This enables a simple and direct selection of suitable mutated operator sequences. A suitable suicide gene is the E lysis gene from the phage PhiX174 as well as homologues and derivatives derived therefrom (Hutchison and Sinsheimer, J.Mol.Biol. 18 (1966), 429-447; Witte et al., Multifunctional safety vector systems for DNA cloning, controlled expression of fusion genes and

- 5 -

simplified preparation of vector DNA and recombinant gene products, in BioTech Forum, Advances in Molecular Genetics 3, pp 219-239, publisher: Issinger, O.-G., Henke, J., Kämpf, J., Driesel, A.J., Hüthing Verlag (1991, Heidelberg). Further examples of suitable lysis genes are GEF (Poulsen et al., Mol.Microbiol. 5 (1991), 1627-1637) and Kil (Reisinger et al., Virology 193 (1993), 1033-1036). On the other hand the selection gene can also be a reporter gene such as e.g. the β -Gal gene.

In order to produce mutants the operator DNA sequence is preferably subjected to a site-specific mutagenesis using one or several oligonucleotides for example according to the method of Kunkel (Proc.Natl.Acad.Sci. USA 82 (1985), 488-492) or they are obtained by selection in a mutator bacterial strain e.g. an E. coli mutD or mutL mutator strain such as E. coli ES1578 (Wu et al., Gene 87 (1990), 1-5). The mutated operator DNA sequences are preferably selected by determining the ability to bind to a temperature-sensitive cI repressor in particular to the temperature-sensitive cI857 repressor. For this the DNA cassette which is preferably located on a vector is transformed into a bacterial cell which contains a gene coding for a temperature-sensitive cI repressor. This gene may also be present on a vector (Remaut et al., Gene 15 (1981), 81-93). On the other hand it is possible to use a bacterial cell which contains such a repressor gene in its chromosome e.g. E. coli M5219 (cf. e.g. Shimatake and Rosenberg, Nature 292 (1981), 128).

Mutants which are resistant to lysis at different temperatures can be identified in a simple manner by culturing the bacterial cells transformed with a lysis cassette which contain the mutated operator DNA

- 6 -

sequences. Up to now it has been possible to identify several mutants which are resistant to a lysis at temperatures up to 33°C, 35°C, 37°C and 39°C. These bacteria contain mutated operator DNA sequences which allow binding of the repressor up to the respective temperature. A particularly preferred example is a mutant to which the cI857 repressor binds up to a temperature of about 37°C. Compared to the wild-type the mutation is a single base substitution in the O_R2 section of the lambda O_R operator region. The sequence of this mutated lambda O_R operator is shown in SEQ ID NO.2.

An additional subject matter of the invention are mutated O_R or O_L operator sequences from lambdoid phages which have a different thermostability compared to the wild-type sequence with regard to binding of a repressor and which are obtainable by the selection methods described above. The mutated O_R or O_L operator sequences preferably have an increased thermostability with regard to the binding of a temperature-sensitive repressor and in particular of the temperature-sensitive cI repressor. The mutated operator sequences particularly preferably have a thermostability that is increased by about 3 to 10°C, in particular by about 7 to 9°C compared to the wild-type sequence.

Since the selection method according to the invention is preferably carried out on O_R or O_L operator sequences which are derived from the phage lambda, the present invention in particular concerns mutated lambda O_R or O_L operator sequences which are variants of the O_R operator sequences shown in SEQ ID NO.1 or variants of the O_L operator sequences shown in SEQ ID NO.3. Variant in this connection is understood as an operator sequence which differs from the wild-type sequence in at least one

- 7 -

sequence position by insertion, deletion or substitution of bases. The differences are particularly preferably in the region of the sections O_R1 , O_R2 or O_R3 and O_L1 , O_L2 and O_L3 . A specific example of a mutated lambda operator sequence according to the invention is the lambda- O_R operator sequence shown in SEQ ID NO.2.

The mutated operator sequences allow the production of new temperature-regulated systems for gene expression in which microorganisms and in particular bacteria can be cultured in a repressed state at variable temperatures and preferably at higher temperatures than have been previously possible in particular at 33 to 39°C. Hence a subject matter of the invention is the use of the mutated O_R or O_L operator sequences for the temperature-regulated expression of genes in bacteria and in particular in gram-negative bacteria such as *E. coli*. Combination of a wild-type O_R or O_L operator region and at least one operator region which contains a mutated operator sequence according to the invention or combination of several operator regions which contain mutated operator sequences according to the invention with different thermostabilities even enables a temperature-regulated sequential expression of genes.

Vectors and bacterial strains in which the inventive mutated operator sequences can be used for the temperature-regulated expression of genes are familiar to a person skilled in the art. In this case the expression systems known from the prior art containing the lambda cI857 repressor in combination with a suitable promoter e.g. the lambda P_L or lambda P_R promoter can be used (cf. e.g. Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd Edition, 1989, Cold Spring Harbor Laboratory Press, New York,

- 8 -

17.11-17.12).

A further subject matter of the present invention is a nucleic acid comprising a bacterial expression control sequence i.e. a sequence containing a promoter and operator regions which contains a mutated O_R or O_L operator sequence according to the invention in operative linkage with a protein-coding sequence. The protein-coding sequence can for example be a sequence coding for a eukaryotic protein or polypeptide or a bacterial gene e.g. the E-lysis gene.

An additional subject matter of the present invention is a vector which contains at least one copy of the bacterial expression control sequence in operative linkage with the protein-coding sequence. This vector can be any prokaryotic vector e.g. a chromosomal vector such as a bacteriophage or an extrachromosomal vector such as a plasmid. Suitable prokaryotic vectors are described for example by Sambrook et al., Supra, chapters 1-4.

Yet a further subject matter of the present invention is a bacterial cell which is transformed with a nucleic acid according to the invention or with a vector according to the invention. In a preferred embodiment the cell is a gram-negative prokaryotic cell, particularly preferably an E. coli cell. The cell preferably contains the nucleic acid or the vector integrated into its chromosome and in addition contains a gene for a cI repressor from a lambdoid phage in particular the gene for the lambda cI_{857} repressor.

A particularly preferred application of the mutated

- 9 -

operators according to the invention is in the field of vaccine production. So-called "bacterial ghosts" are known as vaccines from the prior art i.e. bacterial coats that can be prepared from gram-negative bacteria such as *E. coli*, *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Actinobacillus pleuropneumoniae* etc. by means of protein-E-induced lysis. These ghosts whose cell surface properties and repertoire of surface antigens that can be recognized by the immune system are very similar to the active pathogen, produce a protective cellular or/and humoral immune response in various animal models.

The process for preparing the ghosts is based on the stringent controlled expression of the E-lysis gene from *PhiX174* whose expression product forms a tunnel through the bacterial cell wall coat and thus leads to a pouring out of the cell contents of the host cell. This lethal gene for the cells can be regulated by means of a lambda repressor e.g. the temperature-sensitive lambda repressor *cI857* which, as described above, loses its function at temperatures above 30°C. As a result, the bacterial cultures that have previously been used to produce bacterial ghosts have had to be cultured at low temperatures, preferably at 28°C.

Although this method leads to satisfactory results with regard to the immunogenicity of the ghosts that are produced, an improvement of the bacterial culture is urgently required since the repertoire of antigenic determinants on the bacterial surface can change depending on the external conditions. Since pathogenic bacteria which infect humans or animals usually colonize at an environmental temperature of 37 to 39°C, this natural environmental temperature should also be

- 10 -

maintained during the production process for the ghosts.

A process for producing bacterial ghosts which achieves this object is provided by using the mutated operator sequences according to the invention. These operator sequences allow growth of the bacteria up to a temperature range of preferably 35 to 39°C and allow lysis when the temperature is increased from 37 to 42°C. This changed lysis behaviour enables the pathogens to be cultured near to the body temperature of the vaccine candidate which is extremely important for the composition of the external membrane. Furthermore the new lysis cassette can also be used as a safety cassette in live vaccines since for example in humans the inoculated bacteria are killed when fever is induced (39°C).

Hence a subject matter of the invention is a vaccine composition which contains a live bacterial cell according to the invention as the active ingredient optionally together with pharmaceutically tolerated auxiliary substances, additives and carrier substances. The live bacterial cell contains a nucleic acid comprising a bacterial expression control sequence with a mutated operator sequence in operative linkage preferably with a lysis gene. Yet a further subject matter of the present invention is a vaccine composition which contains a bacterial ghost as an active ingredient optionally together with pharmaceutically tolerated auxiliary substances, additives and carrier substances wherein the bacterial ghost is obtainable by culturing a bacterial cell according to the invention at temperatures of 35 - 39°C and subsequently lysing the bacterial cell by increasing the temperature. Bacterial cells suitable as vaccines are in particular gram-

- 11 -

negative bacteria such as *E. coli* for example the strains STEC, EHEC, O78:K80, *Salmonellae* such as *S. choleraesuis*, *S. enteritidis* and *S. typhimurium*, *Pasteurella multocida*, *Pasteurella haemolytica*, *Bordetella bronchiseptica*, *Klebsiella pneumoniae*, *Actinobacillus pleuropneumoniae*, *Haemophilus influenzae*, *Vibrio cholerae*, *Helicobacter pylori*, *Alcaligenes eutrophus*, *Campylobacter jejuni* and *Pseudomonas aeruginosa*.

The vaccine compositions modified according to the invention can be transferred orally, aerogenically or parenterally to the vaccine candidates. The route which the corresponding microorganisms naturally select for the infection and for the initial stages of establishing an infectious disease are preferably selected for the application of the vaccine. Since all surface properties are retained in the vaccines according to the invention, this application can result in a local induction of the immune response as also occurs in the natural infection process.

As described above the use of mutated operator sequences according to the invention enables the development of vaccines that can be lysed in a controlled manner when a target temperature is exceeded. Furthermore it is, however, also possible to provide a cold-sensitive suicide cassette which on release into the environment kills gram-negative bacteria that are used as a live vaccine. Hence combination of two genetic regulation systems enables the bacteria to die as a result of the expression of a suicide gene when a target value for the environmental temperature is exceeded. This safety cassette ensures that the live vaccines are killed even if they are eliminated from the organisms.

- 12 -

Hence the invention concerns a nucleic acid comprising (a) a first bacterial expression control sequence which contains an O_R or O_L operator sequence from a lambdoid phage and to which a first temperature-sensitive cI repressor from lambdoid phages can bind in operative linkage with a sequence coding for a second repressor wherein the second repressor cannot bind to the first bacterial expression sequence and (b) a second bacterial expression control sequence to which the second repressor can bind which is in operative linkage with a suicide gene.

The components (a) and (b) can be covalently linked together e.g. be present on a single vector or be separate from one another e.g. present on different vectors or be located separately or together on the chromosome of a recipient bacterium.

Yet a further subject matter of the present invention is a bacterial cell which contains at least one copy of a nucleic acid as described above. In addition the bacterial cell advantageously contains a gene for the first repressor. The first repressor is preferably the temperature-sensitive cI857 repressor.

The safety cassette according to the invention preferably contains a gene which codes for a temperature-sensitive cI repressor e.g. the repressor cI857 and a gene which codes for a second repressor wherein this gene is under the control of a lambda promoter/operator region to which the temperature-sensitive repressor binds. The second repressor in turn controls the expression of another gene e.g. a suicide gene such as the E-lysis gene. The temperature-sensitive

- 13 -

lambda repressor is inactive at 37°C so that the second repressor is expressed which in turn represses the expression of the suicide gene.

When the temperature is reduced the temperature-sensitive lambda repressor binds to the operator so that the expression of the second repressor is blocked which leads to an expression of the suicide gene. A first expression control sequence is preferred for this safety cassette which contains the mutated lambda operator since this enables an improved and more rapid activation of the suicide gene.

The second repressor can be any repressor e.g. a lac repressor. However, it is preferable to use an additional repressor from lambdoid phages e.g. cI from the phage 434 which is not temperature-sensitive and binds to its own operator sequence but does not bind to the sequence recognized by the lambda repressor cI857.

It is particularly preferable for the development of live vaccines to incorporate a heat as well as a cold regulation element. This incorporation is preferably achieved by homologous recombination into the chromosome of the vaccine bacterium.

Thus the present invention also concerns a bacterial cell which, in addition to the two components (a) and (b), contains a third bacterial expression control sequence as component (c) which contains a mutated operator sequence according to the invention in operative linkage with a suicide gene.

- 14 -

These bacterial cells can also be used in vaccine compositions and especially for live vaccines. In this manner it is possible to produce heat or/and cold-sensitive safe live vaccines which lead to death of the vaccine bacteria when the body temperature of the vaccine candidate is increased e.g. by fever or/and when they are excreted into the environment.

It is intended to additionally elucidate the invention by the following figures, sequence protocols and examples.

- Fig. 1a shows a schematic representation of a lysis cassette of the prior art comprising a lambda O_R wild-type region, the lambda $cI857$ gene under the control of the promoter P_{RM} and the E lysis gene under the control of the promoter P_R ;
- Fig. 1b shows a schematic representation of a lysis cassette according to the invention which contains a mutated lambda O_R sequence;
- Fig. 2a shows a schematic representation of a cold-sensitive safety cassette comprising a wild-type (pCS1) or mutated (pCSJ1) O_R operator sequence, the lambda- $cI857$ gene under the control of the promoter P_{RM} , the gene of the $lacI$ repressor under the control of P_R and the E-lysis gene under the control of the lac promoter/operator system at a temperature at which the temperature-sensitive lambda repressor $cI857$ does not bind to the lambda O_R sequence;
- Fig. 2b shows a schematic representation of the safety cassette according to Fig. 2a at a temperature at which the lambda repressor $cI857$ binds to the lambda O_R operator;

- 15 -

- Fig. 3 shows the lysis curve of bacterial cells (optical density versus time) which contain a plasmid with the lysis cassette shown in Fig. 1b;
- Fig. 4 shows the lysis curve of a bacterial cell which contains a cold-sensitive safety cassette with the wild-type O_R operator and
- Fig. 5 shows a comparison of lysis curves of bacterial cells which contain a cold-sensitive safety lysis cassette with the wild-type O_R operator (pSC1) or the mutated operator (pCSJ1);
- Fig. 6a shows a schematic representation of a cold-sensitive safety cassette comprising a wild-type (pCS2) or mutated (pCSJ2) O_R operator sequence, the lambda cI857 gene under the control of the promoter P_{RM} , the gene of the phage 434 cI repressor under the control of lambda P_R and the E lysis gene under the control of the 434 O_R (P_{RM} - P_R) promoter/operator system at a temperature at which the temperature-sensitive lambda repressor cI857 does not bind to the lambda O_R sequence,
- Fig. 6b shows a schematic representation of the safety cassette according to Fig. 6a at a temperature at which the lambda repressor cI857 binds to the lambda O_R operator;

SEQ ID NO.1 shows the nucleotide sequence of the lambda O_R operator; the operator sequence O_{R3} extends from position 11 - 27; the operator sequence O_{R2} extends from position 34 - 41; the operator sequence O_{R1} extends from position 58 - 74;

SEQ ID NO.2 shows the nucleotide sequence of a mutated lambda O_R operator which, compared to the wild-type sequence, has a substitution of

- 16 -

T → C at position 42;

SEQ ID NO.3 shows the nucleotide sequence of the lambda O_L operator; the operator sequence O_L3 extends from position 11 - 27; the operator sequence O_L2 extends from position 31-- 47; the operator sequence O_L1 extends from position 55 - 70:

SEQ ID NO. 4 to 6

show a 1601 bp long DNA fragment of the plasmid pAW12; bp 1 - 983 are derived from the bacteriophage lambda (position 37125 - 38107; cf. Sanger et al., J.Mol.Biol. 162 (1982), 729-773) and contain the lambda $cI857$ gene (position 816-106; SEQ ID NO.5) as well as the mutated O_R operator region (mutation at position 858 T → C); bp 1023 - 1601 are derived from the phage Φ X174 (position 447 - 1026; cf. Sanger et al., J.Mol.Biol. 125 (1978), 225-246) and contain the E-lysis gene (position 1144 - 1416; SEQ ID NO.6);

SEQ ID NO. 7 to 10

shows a 2834 bp long DNA fragment of the plasmid pCSJ; bp 1 - 983 are derived from the bacteriophage lambda (position 37125 - 38107) and contain the $cI857$ gene (position 816 - 106; SEQ ID NO.5) as well as the mutated lambda O_R region (mutation at position 858 T → C; bp 990 - 2230 are derived from the E. coli lac operon subcloned on the plasmid pMC7 (Calos, Nature 274 (1978), 762-765) and contain the lacI repressor gene (bp 1025 - 2104; SEQ ID NO.9) and the lac promoter/operator; bp 2256 - 2834 are derived from the bacteriophage

- 17 -

PhiX174 (position 447 - 1026) and contain the E-lysis gene (bp 2377 - 2649; SEQ ID NO.10).

Examples

Example 1:

1.1 Random mutagenesis of the lambda O_R operator region

The plasmid pAW12 (Witte and Lubitz, Eur.J.Biochem. 180 (1989), 393-398) was selected as the starting material which contains the lysis gene E from the bacteriophage PhiX174 under the control of the lambda P_R promoter as well as the associated repressor gene cI857. The aim of this experiment was to change the lysis cassette so that the lysis gene E is not already activated at 30°C but at higher temperatures. For this the E. coli mutator strain ES1578 (Wu et al., (1990), supra) was transformed with the lysis plasmid and a selection was carried out for clones with a changed temperature profile of cell lysis.

For this the mutated clones produced by the transformation were detected after being stamped onto test plates containing lysis selective medium (LB containing 1 % SDS) and incubated at different temperatures (e.g. 33°C, 34°C, 35°C, 36°C, 37°C, 38°C, 39°C, 40°C, 41°C). The changed lysis profile of the lysis cassette in liquid culture was exactly determined by plasmid extraction and subsequent transformation into a non-mutator test strain.

The type of mutation was determined by subcloning the

- 18 -

mutagenized lysis cassettes into a sequencing plasmid. In addition the lysis gene E was substituted by the β -galactosidase gene for a functional check. It was then possible on the basis of a simple β -gal test to quantitatively measure the repressed or active state of the gene cassette.

In this manner it was possible to obtain several clones with a different temperature lysis profile. These clones allowed growth of the bacteria in a temperature range of 33-39°C and did not lead to lysis of the bacteria until the temperature was further increased to 37-42°C.

A mutation of the O_R operator region (SEQ ID NO.2) was identified by sequencing a mutated clone which had a thermostability up to 37°C.

1.2 Verification of the mutation

In order to verify the mutation obtained in example 1.1. a site-specific mutagenesis of the lambda O_R wild-type sequence was carried out using an oligonucleotide.

The mutagenesis was carried out according to the protocol of Kunkel (Proc.Natl.Acad.Sci. USA 82 (1985), 488-492).

4 ml overnight culture of the E. coli strain CJ236 (dut⁻, ung⁻) was added to 50 ml LB medium (+ 10 μ g/ml chloramphenicol and 0.25 μ g/ml uridine) and shaken for 30 min at 37°C. Then 100 μ l M13 phages was added and it was incubated for 6 h at 37°C.

- 19 -

The culture was centrifuged in 2 SS34 centrifuge tubes for 10 min at 14000 rpm and 4°C, the supernatant was decanted into new tubes and again centrifuged for further purification.

The phages were precipitated for 1 h at 4°C by addition of 5 ml 5 x polyethylene glycol/NaCl. They were then centrifuged for 10 min at 14000 rpm and 4°C and the supernatant was discarded.

The pellet was dried, suspended in 0.8 ml TES buffer (0.1 M Tris HCl, pH 8; 0.3 M NaCl; 1 mM EDTA) and incubated for 1 h at 4°C. The suspension was divided into 2 Eppendorf vessels and centrifuged for 5 min at 5000 rpm. The supernatant in which the disrupted phages were located was removed and subjected to a phenol/chloroform extraction to isolate the DNA. The resulting DNA was precipitated with a 2.5-fold volume of 96 % ethanol, washed with 70 % ethanol and taken up in 60 µl H₂O.

An oligonucleotide with the sequence
5'-GTA AAA TAG TCA ACA CGC GCG GTG TTA GAT ATT TAT C-3'
was phosphorylated. For this 20 µl H₂O, 20 µl oligonucleotide (20 ng), 4.5 µl kinase buffer (Stratagene) and 0.5 µl polynucleotide kinase (5 U, Stratagene) was incubated for 1 h at 37°C. Then 7 µl 0.1 M EDTA was added and it was heated for 10 min to 65°C.

For the annealing 20 µl phosphorylated oligonucleotide, 3.5 µl single-stranded DNA template (1 µg ssDNA produced as described above) and 1.4 µl 20 x SSC buffer were heated for 5 min to 70°C, slowly cooled to 25°C and then

- 20 -

placed on ice.

For the extension 10 μ l of the reaction mixture from the annealing mixture, 37.5 μ l XL buffer (27 mM Hepes pH 7.8, 5 mM of each dNTP, 13 mM $MgCl_2$, 2.7 mM dithiothreitol, 1.3 mM ATP, 1 μ l ligase (1 U, Boehringer Mannheim), 1.5 μ l T4 polymerase (1.5 U, Boehringer Mannheim), 1.5 μ l T4 gene32 protein (8 μ g, Boehringer Mannheim) were incubated for 10 min on ice, 10 min at room temperature and 2 h at 37°C. After 1 h 1 μ l ligase and 1 μ l T4 DNA polymerase was added. After completion of the incubation the reaction was stopped by adding 3 μ l 0.25 M EDTA.

For the transformation 100 μ l competent E. coli cells JM103 (Messing et al., Nucleic Acids Res. 9 (1981), 309-321) was admixed with 10 μ l DNA from the extension mixture and incubated for 1 h or more on ice. After a heat shock for 2.5 min at 42°C, 0.2 ml fresh JM103 cells was added in the logarithmic growth phase. The cells were mixed with 3 ml soft agar and inoculated on an LB agar plate. They were subsequently incubated overnight at 37°C.

In order to identify the mutants, plaques were pricked out with a Pasteur pipette and used to inoculate 5 ml LB medium to which 400 μ l of an overnight culture of E. coli JM103 had been added. After 3 h growth at 37°C, the cells were centrifuged. Double-stranded M13 plasmids were obtained from the cell pellet by means of plasmid preparation. Single-stranded M13 phages could be isolated from the supernatant.

- 21 -

Example 2:Analysis of the mutagenized lysis cassettes

Figures 1 and 2 shows different E-specific lysis cassettes with different temperature inductions of the lysis function.

In Fig. 1a which contains the wild-type lambda O_R operator sequence, the function of the E-lysis gene is repressed up to 30°C by the cI857-coded repressor protein on the preceding lambda P_R promoter/operator region. cI857-specific repressor molecules are thermally inactivated at temperatures above 30°C and the expression of the E gene is induced. Fig. 1b shows the plasmid pAWJ12 which contains a mutated operator sequence (SEQ ID NO.2) so that the repression of the lysis function of the gene E by cI857 occurs up to 37°C and the lysis function is not induced until 39°C or higher temperatures are reached.

The function of a cold-sensitive safety cassette is elucidated in Fig. 2. Fig. 2a shows that the formation of lacI-specific repressor molecules which in turn repress the expression of the E gene is induced in the plasmids pCS1 (wild-type operator) and pCSJ1 (mutated operator) at a temperature of $\geq 32^\circ\text{C}$ (pCS1) or $\geq 39^\circ\text{C}$ (pCSJ1). At a temperature below 30°C (pCS1) or 37°C (pCSJ1) functional cI857 repressor molecules are formed which suppress the formation of lacI-specific repressor molecules and thus allow the expression of the E gene (Fig. 2b). In the plasmid pCSJ1 the concomitant cell lysis occurs more rapidly than in pCS1.

- 22 -

Fig. 3 shows the lysis curve of a bacterial cell containing the plasmid pAWJ12 (mutated operator). 3 hours after beginning the culture, the temperature was maintained at 37°C in an aliquot of the bacterial cells and increased in two other aliquots to 38 and 42°C. At 37°C there was a further growth of the bacteria whereas a lysis already occurred at 38°C. The lysis is considerably increased at 42°C.

Figures 4 and 5 show the function of a cold-sensitive safety cassette. In Fig. 4 bacterial cells which contained the plasmid pCS1 (wild-type operator) were subjected to a temperature change from 37 to 28°C. This reduction in temperature led to the E-lysis gene being switched off and cell death (decrease of the optical density).

Fig. 5 shows a comparison of the lysis rate of bacteria which contain the plasmid pCS1 (wild-type operator) and the plasmid pCSJ1 (mutated operator). It can be seen that the lysis occurs much more rapidly in the bacteria which contain the mutated operator.

Fig. 6 shows a further cold-sensitive safety cassette. At temperatures at which the lambda cI857 repressor does not bind to the operator the plasmids pCS2 (wild-type operator) and pCSJ2 (mutated operator) form cI-434 repressor molecules which repress the expression of the E gene (Fig. 6a). Formation of cI-434-specific repressor molecules is prevented thus allowing expression of the E gene at a temperature at which the cI857 repressor binds to the lambda operator (Fig. 6b).

- 23 -

Example 3:In vivo analysis of cold-sensitive lysis cassettes

The killing of bacteria by lowering the temperature after passage through a mouse intestine and excretion into the faeces was determined.

For this 10^{10} E. coli bacteria were administered once to Balb/c mice and the excreted number of bacteria in the faeces was determined. The evaluation was carried out on E.coli-specific Endo plates (Endo, "Zentralbl. Bakt. I Orig." 35 (1904) 109-110) using tetracyclin as a marker for the plasmids used. The incubation was carried out at 28°C.

Results:

In the experimental groups E. coli NM522 (pCS2), E. coli MC4100 (pCS1) and E. coli MC4100 (pCSJ1) there was a reduction in the germ count compared to an E. coli NM522 (pAWJ-lac) control of at least 99.9 %, 98 % and 80 % measured 10 h and 20 h after administering the E. coli bacteria.

- 24 -

SEQUENCE PROTOCOL

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Prof.DR. Werner Lubitz
- (B) ROAD: Schoenborngasse 12/7
- (C) CITY: Vienna
- (E) COUNTRY: Austria
- (F) POSTAL CODE: 1080

(ii) TITLE OF INVENTION: New systems for the regulation
of gene expression

(iii) NUMBER OF SEQUENCES: 10

(iv) COMPUTER-READABLE FORM:

- (A) DATA CARRIER: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0,
Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 82 base pairs
- (B) TYPE: nucleotide
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: linear

(vi) INITIAL ORIGIN:

- (A) ORGANISM: lambda OR operator (wild-type)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
ACGTTAAATC TATCACCGCA AGGATAAAT ATCTAACACC GTGCGTGTG ACTATTTTAC      60
CTCTGGCGGT GATAATGGTT GC                                             82
```

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 82 base pairs
- (B) TYPE: nucleotide
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: linear

- 25 -

(vi) INITIAL ORIGIN:
 (A) ORGANISM: lambda OR operator (mutant)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

ACGTTAAATC TATCACCGCA AGGGATAAAT ATCTAACACC GCGCGTGTG ACTATTTTAC      60
CTCTGGCGGT GATAATGGTT GC .                                           82

```

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 85 base pairs
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

(vi) INITIAL ORIGIN:
 (A) ORGANISM: lambda OL operator (wild-type)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

```

ACATACAGAT AACCATCTGC GGTGATAAAT TATCTCTGGC GGTGTTGACA TAAATACCAC      60
TGGCGGTGAT ACTGAGCACA TCAGC                                           85

```

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1601 base pairs
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: double strand
 (D) TOPOLOGY: both

(vi) INITIAL ORIGIN:
 (A) ORGANISM: pAW12 fragment

(xi) CHARACTERISTICS:
 (A) NAME/KEY: CDS
 (B) LOCATION: complement (106..816)

- 26 -

(ix) CHARACTERISTICS:

(A) NAME/KEY: CDS

(B) LOCATION: 1144..1416

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

10

15 ATTTACTATG TTATGTTCTG AGGGGAGTGA AAATTCCTCT AATTCGATGA AGATTCTTGC 60
 TCAATTGTGA TCAGCTATGC GCCGACCAGA ACACCTTGCC GATCAGCCAA ACGTCTCTTC 120
 AGGCCACTGA CTAGCGATAA CTTTCCCCAC AACCGAACAA CTCTCATTCG ATGGGATCAT 180
 20 TGGGTACTGT GGGTTTAGTG GTTGTAAGAA CACCTGACCG CTATCCCTGA TCAGTTTCTT 240
 GAAGGTAAAC TCATCACCCC CAAGTCTGGC TATGCAGAAA TCACCTGGCT CAACAGCCTG 300
 CTCAGGGTCA ACGAGAATTA ACATTCCGTC AGGAAAGCTT GGCTTGAGC CTGTTGGTGC 360
 25 GGTCATGGAA TTACCTTCAA CCTCAAGCCA GAATGCAGAA TCACTGGCTT TTTTGGTTGT 420
 GCTTACCCAT CTCTCCGCAT CACCTTGGT AAAGTTCTA AGCTTAGGTG AGAACATCCC 480
 30 TGCCTGAACA TGAGAAAAAA CAGGGTACTC ATACTCACTT CTAAGTGACG GCTGCATACT 540
 AACCCTTCA TACATCTCGT AGATTCTCT GGCATTGAA GGGCTAAAT CTTCAACGCT 600
 AACTTTGAGA ATTTTGTAA GCAATGCGGC GTTATAAGCA TTTAATGCAT TGATGCCATT 660
 35 AAATAAGCA CCAACGCCTG ACTGCCCCAT CCCCCTCTG TCTGCGACAG ATTCCTGGGA 720
 TAAGCCAAGT TCATTTTCT TTTTTCATA AATTGCTTA AGGCGACGTG CGTCCTCAAG 780
 40 CTGCTCTGT GTTAATGGT TCTTTTGT GCTCATACGT TAAATCTATC ACCGCAAGGG 840
 ATAATATCT AACACGCGC GTGTGACTA TTTTACCTCT GCGGTGATA ATGGTTGCAT 900
 GTACTAAGTA GGTGTATGG AACACGCAT AACCTGAAA GATTATGCAA TCGCCTTGG 960
 45 GCAAACCAAG ACAGCTAAAG ATCCTCTAGA GTCGACCTG AGGCATGCAA GCTTATCGAA 1020
 TTCTCATTCA GGCTTCTGCC GTTTTGGATT TAACCGAAGA TGATTTGAT TTTCTGACGA 1080
 50 GTAACAAAGT TTGATTGCT ACTGACCGCT CTCGTGCTCG TCGCTGCGT GAGGCTTGCG 1140
 TTT ATG GTA CGC TGG ACT TTG TGG GAT ACC CTC GCT TTC CTG CTC CTG 1188
 Met Val Arg Trp Thr Leu Trp Asp Thr Leu Ala Phe Leu Leu Leu
 1 5 10 15
 55 TTG AGT TTA TTG CTG CCG TCA TTG CTT ATT ATG TTC ATC CCG TCA ACA 1236
 Leu Ser Leu Leu Leu Pro Ser Leu Leu Ile Met Phe Ile Pro Ser Thr
 20 25 30
 60 TTC AAA CGG CCT GTC TCA TCA TGG AAG GCG CTG AAT TTA CGG AAA ACA 1284
 Phe Lys Arg Pro Val Ser Ser Trp Lys Ala Leu Asn Leu Arg Lys Thr
 35 40 45

65

20

25

25

25

25

25

30

30

35	Met	Ser	Thr	Lys	Lys	Pro	Leu	Thr	Gln	Glu	Gln	Leu	Glu	Asp	Ala
	1				5				10					15	
	Arg	Arg	Leu	Lys	Ala	Ile	Tyr	Glu	Lys	Lys	Lys	Asn	Glu	Leu	Gly
				20					25				30		Leu
40	Ser	Gln	Glu	Ser	Val	Ala	Asp	Lys	Met	Gly	Met	Gly	Gln	Ser	Gly
			35					40					45		Val
	Gly	Ala	Leu	Phe	Asn	Gly	Ile	Asn	Ala	Leu	Asn	Ala	Tyr	Asn	Ala
45		50					55					60			Ala
	Leu	Leu	Thr	Lys	Ile	Leu	Lys	Val	Ser	Val	Glu	Glu	Phe	Ser	Pro
	65					70					75				80
50	Ile	Ala	Arg	Glu	Ile	Tyr	Glu	Met	Tyr	Glu	Ala	Val	Ser	Met	Gln
					85					90					95
	Ser	Leu	Arg	Ser	Glu	Tyr	Glu	Tyr	Pro	Val	Phe	Ser	His	Val	Gln
				100					105					110	Ala
55	Gly	Met	Phe	Ser	Pro	Lys	Leu	Arg	Thr	Phe	Thr	Lys	Gly	Asp	Ala
			115					120					125		Glu
	Arg	Trp	Val	Ser	Thr	Thr	Lys	Lys	Ala	Ser	Asp	Ser	Ala	Phe	Trp
60		130					135					140			Leu
	Glu	Val	Glu	Gly	Asn	Ser	Met	Thr	Ala	Pro	Thr	Gly	Ser	Lys	Pro
	145					150					155				160

- 28 -

Phe Pro Asp Gly Met Leu Ile Leu Val Asp Pro Glu Gln Ala Val Glu
 165 170 175
 Pro Gly Asp Phe Cys Ile Ala Arg Leu Gly Gly Asp Glu Phe Thr Phe
 5 180 185 190
 Lys Lys Leu Ile Arg Asp Ser Gly Gln Val Phe Leu Gln Pro Leu Asn
 195 200 205
 10 Pro Gln Tyr Pro Met Ile Pro Cys Asn Glu Ser Cys Ser Val Val Gly
 210 215 220
 Lys Val Ile Ala Ser Gln Trp Pro Glu Glu Thr Phe Gly
 15 225 230 235

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 91 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) TYPE OF MOLECULE: protein

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

30

Met Val Arg Trp Thr Leu Trp Asp Thr Leu Ala Phe Leu Leu Leu Leu
 1 5 10 15
 35 Ser Leu Leu Leu Pro Ser Leu Leu Ile Met Phe Ile Pro Ser Thr Phe
 20 25 30
 Lys Arg Pro Val Ser Ser Trp Lys Ala Leu Asn Leu Arg Lys Thr Leu
 35 40 45
 40 Leu Met Ala Ser Ser Val Arg Leu Lys Pro Leu Asn Cys Ser Arg Leu
 50 55 60
 45 Pro Cys Val Tyr Ala Gln Glu Thr Leu Thr Phe Leu Leu Thr Gln Lys
 65 70 75 80
 Lys Thr Cys Val Lys Asn Tyr Val Gln Lys Glu
 85 90

50

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- 55 (A) LENGTH: 2834 base pairs
 (B) TYPE: nucleotide
 (C) STRANDEDNESS: double strand
 (D) TOPOLOGY: both

60

- 29 -

(vi) INITIAL ORIGIN:

(A) ORGANISM: pCSJ fragment

(xi) CHARACTERISTICS:

(A) NAME/KEY: CDS

(B) LOCATION: complement (106..816)

(ix) CHARACTERISTICS:

(A) NAME/KEY: CDS

(B) LOCATION: 1025..2104

(ix) CHARACTERISTICS:

(A) NAME/KEY: CDS

(B) LOCATION: 2377..2649

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

20

ATTTACTATG TTATGTTCTG AGGGGAGTGA AAATCCCT AATTCGATGA AGATTCTTGC 60

25 TCAATTGTTA TCAGCTATGC GCCGACCAGA ACACCTTGCC GATCAGCCAA ACGTCTCTTC 120

AGGCCACTGA CTAGCGATAA CTTTCCCCAC AACGGAACAA CTCTCATTGC ATGGGATCAT 180

TGGGTACTGT GGGTTTAGTG GTTGTAAGAA CACCTGACCG CTATCCCTGA TCAGTTTCTT 240

30 GAAGGTAAAC TCATCACCCC CAAGTCTGGC TATGCAGAAA TCACCTGGCT CAACAGCCTG 300

CTCAGGGTCA ACGAGAATTA ACATTCCGTC AGGAAAGCTT GGCTTGGAGC CTGTTGGTGC 360

35 GGTCATGGAA TTACCTTCAA CCTCAAGCCA GAATGCAGAA TCACTGGCTT TTTTGGTTGT 420

GCTTACCCAT CTCTCCGCAT CACCTTTGGT AAAGGTCTA AGCTTAGGTG AGAACATCCC 480

TGCCTGAACA TGAGAAAAAA CAGGGTACTC ATACTCACTT CTAAGTGACG GCTGCATACT 540

40 AACCGCTTCA TACATCTCGT AGATTCTCTT GCGGATTGAA GGGCTAAAT CTTCAACGCT 600

AACTTTGAGA ATTTTGTAA GCAATGCGGC GTTATAAGCA TTTAATGCAT TGATGCCATT 660

45 AAATAAGCA CCAACGCCTG ACTGCCCCAT CCCCATCTTG TCTGCGACAG ATTCCTGGGA 720

TAAGCCAAGT TCATTTTCT TTTTTCATA AATTGCTTTA AGGCGACGTG CGTCCTCAAG 780

CTGCTCTTGT GTTAATGGTT TCTTTTGTG GCTCATACGT TAAATCTATC ACCGCAAGGG 840

50 ATAAATATCT AACACGCGC GTGTTGACTA TTTTACCTCT GCGGGTGATA ATGGTTGCAT 900

GTACTAAGTA GGTGTATGG AACAACGCAT AACCCTGAAA GATTATGCAA TCGCCTTTGG 960

55 GCAAACCAAG ACAGCTAAAG ATCCTCTAGA GCGCCCGGAA GAGAGTCAAT TCAGGGTGGT 1020

GAAT GTG AAA CCA GTA ACG TTA TAC GAT GTC GCA GAG TAT GCC GGT GTC 1069

Val Lys Pro Val Thr Leu Tyr Asp Val Ala Glu Tyr Ala Gly Val

95 100 105

60 TCT TAT CAG ACC GTT TCC CGC GTG GTG AAC CAG GCC AGC CAC GTT TCT 1117

Ser Tyr Gln Thr Val Ser Arg Val Val Asn Gln Ala Ser His Val Ser

110 115 120

- 30 -

	GCG AAA ACG CGG GAA AAA GTG GAA GCG GCG ATG GCG GAG CTG AAT TAC	1165
	Ala Lys Thr Arg Glu Lys Val Glu Ala Ala Met Ala Glu Leu Asn Tyr	
	125 130 135	
5	ATT CCC AAC CGC GTG GCA CAA CAA CTG GCG GGC AAA CAG TCG TTG CTG	1213
	Ile Pro Asn Arg Val Ala Gln Gln Leu Ala Gly Lys Gln Ser Leu Leu	
	140 145 150	
10	ATT GGC GTT GCC ACC TCC AGT CTG GCC CTG CAC GCG CCG TCG CAA ATT	1261
	Ile Gly Val Ala Thr Ser Ser Leu Ala Leu His Ala Pro Ser Gln Ile	
	155 160 165 170	
15	GTC GCG GCG ATT AAA TCT CGC GCC GAT CAA CTG GGT GCC AGC GTG GTG	1309
	Val Ala Ala Ile Lys Ser Arg Ala Asp Gln Leu Gly Ala Ser Val Val	
	175 180 185	
20	GTG TCG ATG GTA GAA CGA AGC GGC GTC GAA GCC TGT AAA GCG GCG GTG	1357
	Val Ser Met Val Glu Arg Ser Gly Val Glu Ala Cys Lys Ala Ala Val	
	190 195 200	
25	CAC AAT CTT CTC GCG CAA CGC GTC AGT GGG CTG ATC ATT AAC TAT CCG	1405
	His Asn Leu Leu Ala Gln Arg Val Ser Gly Leu Ile Ile Asn Tyr Pro	
	205 210 215	
30	CTG GAT GAC CAG GAT GCC ATT GCT GTG GAA GCT GCC TGC ACT AAT GTT	1453
	Leu Asp Asp Gln Asp Ala Ile Ala Val Glu Ala Ala Cys Thr Asn Val	
	220 225 230	
35	CCG GCG TTA TTT CTT GAT GTC TCT GAC CAG ACA CCC ATC AAC AGT ATT	1501
	Pro Ala Leu Phe Leu Asp Val Ser Asp Gln Thr Pro Ile Asn Ser Ile	
	235 240 245 250	
40	ATT TTC TCC CAT GAA GAC GGT ACG CGA CTG GGC GTG GAG CAT CTG GTC	1549
	Ile Phe Ser His Glu Asp Gly Thr Arg Leu Gly Val Glu His Leu Val	
	255 260 265	
45	GCA TTG GGT CAC CAG CAA ATC GCG CTG TTA GCG GGC CCA TTA AGT TCT	1597
	Ala Leu Gly His Gln Gln Ile Ala Leu Leu Ala Gly Pro Leu Ser Ser	
	270 275 280	
50	GTC TCG GCG CGT CTG CGT CTG GCT GGC TGG CAT AAA TAT CTC ACT CGC	1645
	Val Ser Ala Arg Leu Arg Leu Ala Gly Trp His Lys Tyr Leu Thr Arg	
	285 290 295	
55	AAT CAA ATT CAG CCG ATA GCG GAA CGG GAA GGC GAC TGG AGT GCC ATG	1693
	Asn Gln Ile Gln Pro Ile Ala Glu Arg Glu Gly Asp Trp Ser Ala Met	
	300 305 310	
60	TCC GGT TTT CAA CAA ACC ATG CAA ATG CTG AAT GAG GGC ATC GTT CCC	1741
	Ser Gly Phe Gln Gln Thr Met Gln Met Leu Asn Glu Gly Ile Val Pro	
	315 320 325 330	
65	ACT GCG ATG CTG GTT GCC AAC GAT CAG ATG GCG CTG GGC GCA ATG CGC	1789
	Thr Ala Met Leu Val Ala Asn Asp Gln Met Ala Leu Gly Ala Met Arg	
	335 340 345	
70	GCC ATT ACC GAG TCC GGG CTG CGC GTT GGT GCG GAT ATC TCG GTA GTG	1837
	Ala Ile Thr Glu Ser Gly Leu Arg Val Gly Ala Asp Ile Ser Val Val	
	350 355 360	
75	GGA TAC GAC GAT ACC GAA GAC AGC TCA TGT TAT ATC CCG CCG TCA ACC	1885
	Gly Tyr Asp Asp Thr Glu Asp Ser Ser Cys Tyr Ile Pro Pro Ser Thr	
	365 370 375	
80	ACC ATC AAA CAG GAT TTT CGC CTG CTG GGG CAA ACC AGC GTG GAC CGC	1933
	Thr Ile Lys Gln Asp Phe Arg Leu Leu Gly Gln Thr Ser Val Asp Arg	
	380 385 390	

- 31 -

	TTG CTG CAA CTC TCT CAG GGC CAG GCG GTG AAG GGC AAT CAG CTG TTG	1981
	Leu Leu Gln Leu Ser Gln Gly Gln Ala Val Lys Gly Asn Gln Leu Leu	
	395 400 405 410	
5	CCC GTC TCA CTG GTG AAA AGA AAA ACC ACC CTG GCG CCC AAT ACG CAA	2029
	Pro Val Ser Leu Val Lys Arg Lys Thr Thr Leu Ala Pro Asn Thr Gln	
	415 420 425	
10	ACC GCC TCT CCC CGC GCG TTG GCC GAT TCA TTA ATG CAG CTG GCA CGA	2077
	Thr Ala Ser Pro Arg Ala Leu Ala Asp Ser Leu Met Gln Leu Ala Arg	
	430 435 440	
15	CAG GTT TCC CGA CTG GAA AGC GGG CAG TGAGCGCAAC GCAATTAATG	2124
	Gln Val Ser Arg Leu Glu Ser Gly Gln	
	445 450	
	TGAGTTAGCT CACTCATTAG GCACCCCAAG CTTTACACTT TATGCTTCCG GCTCGTATGT	2184
20	TGTGTGGAAT TGTGAGCGGA TAACAATTTC ACACAGGAAA CAGCTCTGCA GGCATGCAAG	2244
	CTTATCGAAT TCTCATTCAG GCTTCTGCCG TTTTGGATT T AACCGAAGAT GATTTCGATT	2304
	TTCTGACGAG TAACAAAGTT TGGATTGCTA CTGACCGCTC TCGTGCTCGT CGCTGCGTTG	2364
25	AGGCTTGCGT TT ATG GTA CGC TGG ACT TTG TGG GAT ACC CTC GCT TTC	2412
	Met Val Arg Trp Thr Leu Trp Asp Thr Leu Ala Phe	
	1 5 10	
30	CTG CTC CTG TTG AGT TTA TTG CTG CCG TCA TTG CTT ATT ATG TTC ATC	2460
	Leu Leu Leu Leu Ser Leu Leu Leu Pro Ser Leu Leu Ile Met Phe Ile	
	15 20 25	
35	CCG TCA ACA TTC AAA CGG CCT GTC TCA TCA TGG AAG GCG CTG AAT TTA	2508
	Pro Ser Thr Phe Lys Arg Pro Val Ser Ser Trp Lys Ala Leu Asn Leu	
	30 35 40	
40	CGG AAA ACA TTA TTA ATG GCG TCG AGC GTC CGG TTA AAG CCG CTG AAT	2556
	Arg Lys Thr Leu Leu Met Ala Ser Ser Val Arg Leu Lys Pro Leu Asn	
	45 50 55 60	
45	TGT TCG CGT TTA CCT TGC GTG TAC GCG CAG GAA ACA CTG ACG TTC TTA	2604
	Cys Ser Arg Leu Pro Cys Val Tyr Ala Gln Glu Thr Leu Thr Phe Leu	
	65 70 75	
50	CTG ACG CAG AAG AAA ACG TGC GTC AAA AAT TAC GTG CAG AAG GAG	2649
	Leu Thr Gln Lys Lys Thr Cys Val Lys Asn Tyr Val Gln Lys Glu	
	80 85 90	
55	TGATGTAATG TCTAAAGGTA AAAACGTTT TGGCGCTCGC CCTGGTCGTC CGCAGCCGTT	2709
	GCGAGGTACT AAAGGCAAGC GTAAAGGCGC TCGTCTTTGG TATGTAGGTG GTCAACAATT	2769
	TTAATTGCAG GGGCTTCGGC CCTTACTTGA GGATAAATTA TGTCTAATAT TCAAACCTGGC	2829
	GCCGA	2834

60 (2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- 65 (A) LENGTH: 237 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

- 32 -

(ii) TYPE OF MOLECULE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

5 Met Ser Thr Lys Lys Lys Pro Leu Thr Gln Glu Gln Leu Glu Asp Ala
  1      5      10      15
Arg Arg Leu Lys Ala Ile Tyr Glu Lys Lys Lys Asn Glu Leu Gly Leu
  20      25      30
10 Ser Gln Glu Ser Val Ala Asp Lys Met Gly Met Gly Gln Ser Gly Val
  35      40      45
15 Gly Ala Leu Phe Asn Gly Ile Asn Ala Leu Asn Ala Tyr Asn Ala Ala
  50      55      60
Leu Leu Thr Lys Ile Leu Lys Val Ser Val Glu Glu Phe Ser Pro Ser
  65      70      75      80
20 Ile Ala Arg Glu Ile Tyr Glu Met Tyr Glu Ala Val Ser Met Gln Pro
  85      90      95
Ser Leu Arg Ser Glu Tyr Glu Tyr Pro Val Phe Ser His Val Gln Ala
  100      105      110
25 Gly Met Phe Ser Pro Lys Leu Arg Thr Phe Thr Lys Gly Asp Ala Glu
  115      120      125
Arg Trp Val Ser Thr Thr Lys Lys Ala Ser Asp Ser Ala Phe Trp Leu
  130      135      140
Glu Val Glu Gly Asn Ser Met Thr Ala Pro Thr Gly Ser Lys Pro Ser
  145      150      155      160
35 Phe Pro Asp Gly Met Leu Ile Leu Val Asp Pro Glu Gln Ala Val Glu
  165      170      175
Pro Gly Asp Phe Cys Ile Ala Arg Leu Gly Gly Asp Glu Phe Thr Phe
  180      185      190
40 Lys Lys Leu Ile Arg Asp Ser Gly Gln Val Phe Leu Gln Pro Leu Asn
  195      200      205
Pro Gln Tyr Pro Met Ile Pro Cys Asn Glu Ser Cys Ser Val Val Gly
  210      215      220
45 Lys Val Ile Ala Ser Gln Trp Pro Glu Glu Thr Phe Gly
  225      230      235
50

```

(2) INFORMATION FOR SEQ ID NO: 9:

```

55 (i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 360 amino acids
    (B) TYPE: amino acid
    (D) TOPOLOGY: linear
60 (ii) TYPE OF MOLECULE: protein

```

- 33 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

5 Val Lys Pro Val Thr Leu Tyr Asp Val Ala Glu Tyr Ala Gly Val Ser
 1 5 10 15
 Tyr Gln Thr Val Ser Arg Val Val Asn Gln Ala Ser His Val Ser Ala
 20 25 30
 10 Lys Thr Arg Glu Lys Val Glu Ala Ala Met Ala Glu Leu Asn Tyr Ile
 35 40 45
 Pro Asn Arg Val Ala Gln Gln Leu Ala Gly Lys Gln Ser Leu Leu Ile
 50 55 60
 15 Gly Val Ala Thr Ser Ser Leu Ala Leu His Ala Pro Ser Gln Ile Val
 65 70 75 80
 Ala Ala Ile Lys Ser Arg Ala Asp Gln Leu Gly Ala Ser Val Val Val
 85 90 95
 20 Ser Met Val Glu Arg Ser Gly Val Glu Ala Cys Lys Ala Ala Val His
 100 105 110
 25 Asn Leu Leu Ala Gln Arg Val Ser Gly Leu Ile Ile Asn Tyr Pro Leu
 115 120 125
 Asp Asp Gln Asp Ala Ile Ala Val Glu Ala Ala Cys Thr Asn Val Pro
 130 135 140
 30 Ala Leu Phe Leu Asp Val Ser Asp Gln Thr Pro Ile Asn Ser Ile Ile
 145 150 155 160
 Phe Ser His Glu Asp Gly Thr Arg Leu Gly Val Glu His Leu Val Ala
 165 170 175
 35 Leu Gly His Gln Gln Ile Ala Leu Leu Ala Gly Pro Leu Ser Ser Val
 180 185 190
 40 Ser Ala Arg Leu Arg Leu Ala Gly Trp His Lys Tyr Leu Thr Arg Asn
 195 200 205
 Gln Ile Gln Pro Ile Ala Glu Arg Glu Gly Asp Trp Ser Ala Met Ser
 210 215 220
 45 Gly Phe Gln Gln Thr Met Gln Met Leu Asn Glu Gly Ile Val Pro Thr
 225 230 235 240
 Ala Met Leu Val Ala Asn Asp Gln Met Ala Leu Gly Ala Met Arg Ala
 245 250 255
 50 Ile Thr Glu Ser Gly Leu Arg Val Gly Ala Asp Ile Ser Val Val Gly
 260 265 270
 55 Tyr Asp Asp Thr Glu Asp Ser Ser Cys Tyr Ile Pro Pro Ser Thr Thr
 275 280 285
 Ile Lys Gln Asp Phe Arg Leu Leu Gly Gln Thr Ser Val Asp Arg Leu
 290 295 300
 60 Leu Gln Leu Ser Gln Gly Gln Ala Val Lys Gly Asn Gln Leu Leu Pro
 305 310 315 320
 Val Ser Leu Val Lys Arg Lys Thr Thr Leu Ala Pro Asn Thr Gln Thr
 325 330 335
 65

Val Ser Arg Leu Glu Ser Gly Gln
355 360

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 91 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

	Met	Val	Arg	Trp	Thr	Leu	Trp	Asp	Thr	Leu	Ala	Phe	Leu	Leu	Leu	Leu
	1					5					10					15
25	Ser	Leu	Leu	Leu	Pro	Ser	Leu	Leu	Ile	Met	Phe	Ile	Pro	Ser	Thr	Phe
				20					25					30		
	Lys	Arg	Pro	Val	Ser	Ser	Trp	Lys	Ala	Leu	Asn	Leu	Arg	Lys	Thr	Leu
30			35					40					45			
	Leu	Met	Ala	Ser	Ser	Val	Arg	Leu	Lys	Pro	Leu	Asn	Cys	Ser	Arg	Leu
		50					55						60			
35	Pro	Cys	Val	Tyr	Ala	Gln	Glu	Thr	Leu	Thr	Phe	Leu	Leu	Thr	Gln	Lys
	65					70					75					80
	Lys	Thr	Cys	Val	Lys	Asn	Tyr	Val	Gln	Lys	Glu					
					85						90					

- 35 -

Claims

1. Method for selecting mutated O_R or O_L operator DNA sequences from lambdoid phages which have a different thermostability compared to the wild-type sequence with regard to binding a repressor,
wherein
 - (a) a DNA cassette is prepared which contains a selection gene under the operative control of an expression control sequence comprising at least one O_R or O_L operator sequence from a lambdoid phage and a promoter,
 - (b) the operator DNA sequence is subjected to a mutagenesis and
 - (c) the mutated operator DNA sequences are analysed.
2. Method as claimed in claim 1,
wherein
the lambdoid phages are selected from the group comprising the phage lambda, phage 21, phage 22, phage 82, phage 424, phage 434, phage D326, phage DLP12, phage gamma, phage HK022, phage P4, phage Phi80, phage Phi81, coliphage 186 and recombinant variants thereof.
3. Method as claimed in claim 2,
wherein
the phage lambda or recombinant variants thereof are used.

- 36 -

4. Method as claimed in claim 3,
wherein
an operator DNA sequence from the operator regions O_R or/and O_L of the phage lambda is used.
5. Method as claimed in one of the claims 1 - 4,
wherein
the E-lysis gene from the phage PhiX174 is used as the selection gene.
6. Method as claimed in one of the claims 1 - 5,
wherein
the operator DNA sequence is subjected to a site-specific mutagenesis by oligonucleotides or a selection is carried out in a mutator bacterial strain.
7. Method as claimed in one of the claims 1 - 6,
wherein
the mutated operator DNA sequences are analysed by determining their ability to bind to a temperature-sensitive cI repressor.
8. Method as claimed in claim 7,
wherein
the temperature-sensitive lambda repressor cI857 is used.
9. Mutated O_R or O_L operator sequences from lambdoid phages which have a different thermostability compared to the wild-type sequence with regard to binding of a repressor and are obtainable by a method as claimed in one of the claims 1 - 8.

- 37 -

10. Mutated O_R or O_L operator sequences from lambdoid phages which have an increased thermostability compared to the wild-type sequence with regard to binding of a temperature-sensitive repressor and are obtainable by a method as claimed in one of the claims 1 - 8.
11. Mutated O_R or O_L operator sequence as claimed in claim 10,
wherein
it has an approximately 3 - 10°C increased thermostability.
12. Mutated O_R or O_L operator sequence as claimed in claim 10,
wherein
it has an approximately 7 - 9°C increased thermostability.
13. Mutated lambda O_R or O_L operator sequence as claimed in one of the claims 9 - 12, which is a variant of the sequences shown in SEQ ID NO.1 or SEQ ID NO.3.
14. Mutated lambda O_R operator sequence comprising the sequence shown in SEQ ID NO.2.
15. Use of a mutated O_R or O_L operator sequence as claimed in one of the claims 9 - 14 for the temperature-regulated expression of genes in bacterial cells.

- 38 -

16. Use of a combination of (a) a wild-type O_R or O_L operator region and at least one operator region which contains a mutated O_R or O_L operator sequence as claimed in one of the claims 9 - 14 or (b) several operator regions which contain mutated O_R or O_L operator sequences as claimed in one of the claims 9 - 14 with different thermostabilities for the temperature-regulated sequential expression of genes.
17. Use as claimed in claim 15 or 16,
wherein
the bacterial cells contain a gene for a cI repressor from lambdoid phages for the regulation of gene expression.
18. Use as claimed in claim 17,
wherein
the bacterial cells contain the gene for the lambda cI857 repressor.
19. Nucleic acid comprising a bacterial expression control sequence which contains a mutated O_R or O_L operator sequence as claimed in one of the claims 9 - 14 in operative linkage with a protein-coding sequence.
20. Nucleic acid as claimed in claim 19,
wherein
the protein-coding sequence is a suicide gene.
21. Nucleic acid as claimed in claim 20,
wherein
the expression control sequence contains a lambda P_L or P_R promoter.

- 39 -

22. Vector,
wherein
it contains at least one copy of a nucleic acid as
claimed in one of the claims 19 - 21.
23. Vector as claimed in claim 22,
wherein
it is a bacterial chromosomal vector.
24. Vector as claimed in claim 22,
wherein
it is a bacterial extrachromosomal plasmid.
25. Bacterial cell,
wherein
it is transformed with a nucleic acid as claimed in
one of the claims 19 - 21 or with a vector as
claimed in one of the claims 22 - 24.
26. Bacterial cell as claimed in claim 25,
wherein
it contains the nucleic acid or the vector
integrated into its chromosome.
27. Bacterial cell as claimed in claim 25 or 26,
wherein
it additionally contains a gene for a cI repressor
from lambdoid phages.
28. Bacterial cell as claimed in claim 27,
wherein
it contains the gene for the lambda cI857
repressor.

- 40 -

29. Vaccine composition,

wherein

it contains a live bacterial cell as claimed in one of the claims 26 - 28 as an active ingredient optionally with pharmaceutically acceptable auxiliary substances, additives and carrier substances.

30. Vaccine composition,

wherein

it contains a bacterial ghost as the active ingredient optionally with pharmaceutically acceptable auxiliary substances, additives and carrier substances in which the bacterial ghost can be obtained by culturing a bacterial cell as claimed in one of the claims 25 - 28 at temperatures of 35 - 39°C and subsequently lysing the bacterial cell by increasing the temperature.

31. Nucleic acid comprising (a) a first bacterial expression control sequence which contains an O_R or O_L operator sequence from a lambdoid phage and to which a first cI repressor from lambdoid phages can bind, in operative linkage with a sequence coding for a second repressor wherein the second repressor cannot bind to the first bacterial expression sequence and (b) a second bacterial expression control sequence to which the second repressor can bind in operative linkage with a suicide gene.

32. Bacterial cell,

wherein

it contains at least one copy of a nucleic acid as claimed in claim 31.



- 41 -

33. Bacterial cell as claimed in claim 32,
wherein
it additionally contains a gene for the first repressor.
34. Bacterial cell as claimed in claim 32 or 33,
wherein
it contains the first bacterial expression control sequence of a mutated operator sequence as claimed in one of the claims 9 - 14.
35. Bacterial cell as claimed in one of the claims 32 - 34 additionally comprising (c) a third bacterial expression control sequence which contains a mutated operator sequence as claimed in one of the claims 9 - 14 in operative linkage with a suicide gene.
36. Vaccine composition,
wherein
it contains a live bacterial cell as claimed in one of the claims 32 - 35 as the active ingredient optionally together with pharmaceutically acceptable auxiliary substances, additives and carrier substances.
37. Use of vaccine compositions as claimed in claim 29 or 36 as heat-sensitive or/and cold-sensitive safe live vaccines.


**PCT**WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales BüroINTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

(51) Internationale Patentklassifikation ⁶ : C12N 15/73, 15/10, A61K 39/02 // C 12N 9/36	A3	(11) Internationale Veröffentlichungsnummer: WO 98/07874 (43) Internationales Veröffentlichungsdatum: 26. Februar 1998 (26.02.98)
(21) Internationales Aktenzeichen: PCT/EP97/04560 (22) Internationales Anmeldedatum: 21. August 1997 (21.08.97) (30) Prioritätsdaten: 196 33 698:8 21. August 1996 (21.08.96) DE (71)(72) Anmelder und Erfinder: LUBITZ, Werner [AT/AT]; Schönbornrgasse 12/7, A-1080 Wien (AT). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): JECHLINGER, Wolfgang [AT/AT]; Strozzigasse 38/12, A-1080 Wien (AT). SZOSTAK, Michael [AT/AT]; In den Schnablem 9/3, A-2344 Maria Enzersdorf (AT). (74) Anwälte: WEICKMANN, H. usw.; Kopernikusstrasse 9, D-81679 München (DE).	(81) Bestimmungsstaaten: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO Patent (GH, KE, LS, MW, SD, SZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Veröffentlicht <i>Mit internationalem Recherchenbericht.</i> <i>Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist. Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i> (88) Veröffentlichungsdatum des internationalen Recherchenberichts: 26. März 1998 (26.03.98)	
(54) Title: NEW SYSTEMS FOR REGULATING GENETIC EXPRESSION (54) Bezeichnung: NEUE SYSTEME ZUR REGULATION DER GENEXPRESSION (57) Abstract The present invention concerns a process for selecting new P _R - or P _L -operator sequences of lambdoid phages which, compared to wild-type sequences, have a different thermostability for the binding of a repressor. In addition, the invention discloses new mutated P _R - or P _L - operator sequences and their use for temperature-regulated expression of genes and for producing improved vaccines. (57) Zusammenfassung Die vorliegende Erfindung betrifft ein Verfahren zur Selektion neuer P _R - oder P _L -Operatorsequenzen aus lambdoiden Phagen, die eine im Vergleich zur Wildtypsequenz unterschiedliche Thermostabilität hinsichtlich der Bindung eines Repressors aufweisen. Weiterhin werden neue mutierte P _R - oder P _L -Operatorsequenzen sowie deren Verwendung zur temperaturregulierten Expression von Genen und zur Herstellung verbesserter Impfstoffe offenbart.		

1/5

1a) pAW12 $\leq 30^{\circ}\text{C}$  $\geq 30^{\circ}\text{C}$ b) pAWJ12 $\leq 37^{\circ}\text{C}$  $\geq 39^{\circ}\text{C}$


2a)



pCS1

pCSJ1

2b)



pCS1

pCSJ1

2/5




Fig. 3: Growth of *E. coli* NM522 (pAWJ12) when the temperature is changed from 28°C to higher temperatures (↓)

3/5




Fig. 4: Growth of *E. coli* MC4100 (pCS1) when the temperature is changed from 37°C to 28°C (↓)

4/5




Fig. 5: Growth of *E. coli* MC4100 (pCS1) and MC4100 (pCSJ1) when the temperature is changed from 37°C to 28°C (↓)

